

THE ELECTROMAGNETIC FORM FACTORS OF THE PROTON AND NEUTRON: FUNDAMENTAL INDICATORS OF NUCLEON STRUCTURE

GERALD A. MILLER

*Department of Physics, University of Washington
Seattle, WA 98195-1560*

E-mail: miller@phys.washington.edu

We present a relativistic interpretation for why the proton's G_E/G_M falls and QF_2/F_1 is approximately constant. Reproducing the observed G_E^n then mandates the inclusion of the effects of the pion cloud.

1. Introduction

An alternate title would be “Surprises in the Proton”. This talk owes its existence to the brilliant, precise, stunning and exciting recent experimental work on measuring G_E/G_M (or QF_2/F_1) for the proton and G_E for the neutron. My goal here is to interpret the data. Symmetries including Poincaré invariance and chiral symmetry will be the principal tool I’ll use.

If, a few years ago, one had asked participants at a meeting like this about the Q^2 dependence of the proton's G_E/G_M or QF_2/F_1 . Almost everyone one have answered that for large enough values of Q^2 , G_E/G_M would be flat and QF_2/F_1 would fall with increasing Q^2 . The reason for the latter fall being conservation of hadron helicity. Indeed, the shapes of the curves have been obtained in the new measurements, except for the mis-labeling of the ordinate axes. The expected flatness of G_E/G_M holds for QF_2/F_1 , and the quantity G_E/G_M falls rapidly and linearly with Q^2 . This revolutionary behavior needs to be understood!

2. Outline

I will begin with a brief discussion of Light Front Physics. Stan Brodsky has long been advocating this technique, I have become a convert. Then I will discuss a particular relativistic model of the nucleon, and proceed to apply it. The proton calculations will be discussed first, but recent high

accuracy experiments make it necessary for us to compute observables for the neutron as well.

3. Light Front

Light-front dynamics is a relativistic many-body dynamics in which fields are quantized at a “time” $\tau = x^0 + x^3 \equiv x^+$. The τ -development operator is then given by $P^0 - P^3 \equiv P^-$. These equations show the notation that a four-vector A^μ is expressed as $A^\pm \equiv A^0 \pm A^3$. One quantizes at $x^+ = 0$ which is a light-front, hence the name “light front dynamics”. The canonical spatial variable must be orthogonal to the time variable, and this is given by $x^- = x^0 - x^3$. The canonical momentum is then $P^+ = P^0 + P^3$. The other coordinates are $\mathbf{x}_\perp$ and $\mathbf{P}_\perp$.

The most important consequence of this is that the relation between energy and momentum of a free particle is given by: $p_\mu p^\mu = m^2 = p^+ p^- - p_\perp^2 \rightarrow p^- = \frac{p_\perp^2 + m^2}{p^+}$, a relativistic kinetic energy which does not contain a square root operator. This allows the separation of center of mass and relative coordinates, so that the computed wave functions are frame independent.

The use of the light front is particularly relevant for calculating form factors, which are probability amplitudes for an nucleon to absorb a four momentum q and remain a nucleon. The initial and final nucleons have different total momenta. This means that the final nucleon is a boosted nucleon, with different wave function than the initial nucleon. In general, performing the boost is difficult for large values of $Q^2 = -q^2$. However the light front technique allows one to set up the calculation so that the boosts are independent of interactions. Indeed, the wave functions are functions of relative variables and are independent of frame.

3.1. Definitions

Let us define the basic quantities concerning us here. These are the independent form factors defined by

$$\langle N, \lambda' p' | J^\mu | N, \lambda p \rangle = \bar{u}_{\lambda'}(p') \left[F_1(Q^2) \gamma^\mu + \frac{\kappa F_2(Q^2)}{2M_N} i \sigma^{\mu\nu} (p' - p)_\nu \right] u_\lambda(p). \quad (1)$$

The Sachs form factors are defined by the equations:

$$G_E = F_1 - \frac{Q^2}{4M_N^2} \kappa F_2, \quad G_M = F_1 + \kappa F_2. \quad (2)$$

There is an alternate light front interpretation, based on field theory, in which one uses the “good” component of the current, J^+ , to suppress the

effects of quark-pair terms. Then, using nucleon light-cone spinors:

$$F_1(Q^2) = \frac{1}{2P^+} N, \langle \uparrow | J^+ | N, \uparrow \rangle, Q\kappa F_2(Q^2) = \frac{-2M_N}{2P^+} \langle N, \uparrow | J^+ | N, \downarrow \rangle. \quad (3)$$

4. Why I am Giving This Talk

In 1996, Frank, Jennings & I¹ examined the point-like-configuration idea of Frankfurt & Strikman². We needed to start with a relativistic model of the free nucleon. The resulting form factors are shown in Figs. 10 and 11 of our early paper. The function G_M was constrained³ by experimental data to define the parameters of the model, but we predicted a very strong decrease of G_E/G_M as a function of Q^2 . This decrease has now been measured as a real effect, but the task of explaining its meaning remained relevant. That was the purpose of our second paper⁴ in which imposing Poincaré invariance was shown to lead to substantial violation of the helicity conservation rule as well as an analytic result that the ratio QF_2/F_1 is constant for the Q^2 range of the Jefferson Laboratory experiments. Although the second paper is new, the model is the same. Ralston *et al.*⁵ have been talking about non-conservation of helicity for a long time.

5. Three-Body Variables and Boost

We use light front coordinates for the momentum of each of the i quarks, such that $\mathbf{p}_i = (p_i^+, \mathbf{p}_{i\perp})$, $p^- = (p_\perp^2 + m^2)/p^+$. The total (perp)-momentum is $\mathbf{P} = \mathbf{p}_1 + \mathbf{p}_2 + \mathbf{p}_3$, the plus components of the momenta are denoted as

$$\xi = \frac{p_1^+}{p_1^+ + p_2^+}, \quad \eta = \frac{p_1^+ + p_2^+}{P^+}, \quad (4)$$

and the perpendicular relative coordinates are given by

$$\mathbf{k}_\perp = (1 - \xi)\mathbf{p}_{1\perp} - \xi\mathbf{p}_{2\perp}, \quad \mathbf{K}_\perp = (1 - \eta)(\mathbf{p}_{1\perp} + \mathbf{p}_{2\perp}) - \eta\mathbf{p}_{3\perp}. \quad (5)$$

In the center of mass frame we find:

$$\mathbf{p}_{1\perp} = \mathbf{k}_\perp + \xi\mathbf{K}_\perp, \quad \mathbf{p}_{2\perp} = -\mathbf{k}_\perp + (1 - \xi)\mathbf{K}_\perp, \quad \mathbf{p}_{3\perp} = -\mathbf{K}_\perp. \quad (6)$$

The coordinates $\xi, \eta, \mathbf{k}, \mathbf{K}$ are all relative coordinates so that one obtains a frame independent wave function $\Psi(\mathbf{k}_\perp, \mathbf{K}_\perp, \xi, \eta)$.

Now consider the computation of a form factor, taking quark 3 to be the one struck by the photon. One works in a special set of frames with $q^+ = 0$ and $Q^2 = \mathbf{q}_\perp^2$, so that the value of $1 - \eta$ is not changed by the photon. The coordinate $\mathbf{p}_{3\perp}$ is changed to $\mathbf{p}_{3\perp} + \mathbf{q}_\perp$, so only one relative momentum, $\mathbf{K}_\perp$ is changed:

$$\mathbf{K}'_\perp = (1 - \eta)(\mathbf{p}_{1\perp} + \mathbf{p}_{2\perp}) - \eta(\mathbf{p}_{3\perp} + \mathbf{q}_\perp) = \mathbf{K}_\perp - \eta\mathbf{q}_\perp, \quad \mathbf{k}'_\perp = \mathbf{k}_\perp, \quad (7)$$

The arguments of the spatial wave function are taken as the mass-squared operator for a non-interacting system:

$$M_0^2 \equiv \sum_{i=1,3} p_i^- P^+ - P_\perp^2 = \frac{K_\perp^2}{\eta(1-\eta)} + \frac{k_\perp^2 + m^2}{\eta\xi(1-\xi)} + \frac{m^2}{1-\eta}. \quad (8)$$

This is a relativistic version of the square of a the relative three-momentum. Note that the absorption of a photon changes the value to:

$$M_0'^2 = \frac{(K_\perp - \eta q_\perp)^2}{\eta(1-\eta)} + \frac{k_\perp^2 + m^2}{\eta\xi(1-\xi)} + \frac{m^2}{1-\eta}. \quad (9)$$

6. Wave function

Our wave function is based on symmetries. The wave function is anti-symmetric, a function of relative momenta, independent of reference frame, an eigenstate of the spin operator and rotationally invariant (in a specific well-defined sense). The use of symmetries is manifested in the construction of such wave functions, as originally described by Terent'ev⁶, Coester⁷ and their collaborators. A schematic form of the wave functions is

$$\Psi(p_i) = \Phi(M_0^2)u(p_1)u(p_2)u(p_3)\psi(p_1, p_2, p_3), \quad p_i = \mathbf{p}_i s_i, \tau_i \quad (10)$$

where ψ is a spin-isospin color amplitude factor, the p_i are expressed in terms of relative coordinates, the $u(p_i)$ are ordinary Dirac spinors and Φ is a spatial wave function.

We take the the spatial wave function from Schlumpf³:

$$\Phi(M_0) = \frac{N}{(M_0^2 + \beta^2)^\gamma}, \beta = 0.607 \text{ GeV}, \gamma = 3.5, m = 0.267 \text{ GeV}. \quad (11)$$

The value of γ is chosen that $Q^4 G_M(Q^2)$ is approximately constant for $Q^2 > 4 \text{ GeV}^2$ in accord with experimental data. The parameter β helps govern the values of the perp-momenta allowed by the wave function Φ and is closely related to the rms charge radius, and m is mainly determined by the magnetic moment of the proton.

At this point the wave function and the calculation are completely defined. One could evaluate the form factors as $\langle \Psi | J^+ | \Psi \rangle$ and obtain the previous numerical results¹.

7. Simplify Calculation- Light Cone Spinors

The operator $J^+ \sim \gamma^+$ acts its evaluation is simplified by using light cone spinors. These solutions of the free Dirac equation, related to ordinary Dirac spinors by a unitary transformation, conveniently satisfy:

$$\bar{u}_L(p^+, \mathbf{p}', \lambda') \gamma^+ u_L(p^+, \mathbf{p}, \lambda) = 2\delta_{\lambda\lambda'} p^+. \quad (12)$$

To take advantage of this, re-express the wave function in terms of light-front spinors using the completeness relation: $1 = \sum_{\lambda} u_L(p, \lambda) \bar{u}_L(p, \lambda)$. We then find

$$\Psi(p_i) = u_L(p_1, \lambda_1) u_L(p_2, \lambda_2) u_L(p_3, \lambda_3) \psi_L(p_i, \lambda_i), \quad (13)$$

$$\begin{aligned} \psi_L(p_i, \lambda_i) &\equiv [\bar{u}_L(\mathbf{p}_1, \lambda_1) u(\mathbf{p}_1, s_1)] [\bar{u}_L(\mathbf{p}_2, \lambda_2) u(\mathbf{p}_2, s_2)] \\ &\times [\bar{u}_L(\mathbf{p}_3, \lambda_3) u(\mathbf{p}_3, s_3)] \psi(p_1, p_2, p_3). \end{aligned} \quad (14)$$

This is the very same Ψ as before, it is just that now it is easy to compute the matrix elements of the γ^+ operator.

The unitary transformation is also known as the Melosh rotation. The basic point is that one may evaluate the coefficients in terms of Pauli spinors: $|\lambda_i\rangle, |s_i\rangle$, with $\langle \lambda_i | R_M^\dagger(\mathbf{p}_i) | s_i \rangle \equiv \bar{u}_L(\mathbf{p}_i, \lambda_i) u(\mathbf{p}_i, s_i)$. It is easy to show that

$$\langle \lambda_3 | R_M^\dagger(\mathbf{p}_3) | s_3 \rangle = \langle \lambda_3 | \left[\frac{m + (1 - \eta)M_0 + i\boldsymbol{\sigma} \cdot (\mathbf{n} \times \mathbf{p}_3)}{\sqrt{(m + (1 - \eta)M_0)^2 + p_{3\perp}^2}} \right] | s_3 \rangle. \quad (15)$$

The important effect resides in the term $(\mathbf{n} \times \mathbf{p}_3)$ which originates from the lower components of the Dirac spinors. This large relativistic spin effect can be summarized: the effects of relativity are to replace Pauli spinors by Melosh rotation operators acting on Pauli spinors. Thus

$$|\uparrow \mathbf{p}_i\rangle \equiv R_M^\dagger(\mathbf{p}_i) \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad |\uparrow \mathbf{p}_3\rangle \neq |\uparrow\rangle. \quad (16)$$

8. Proton F_1, F_2 -Analytic Insight

The analytic insight is based on Eq. (15). Consider high momentum transfer such that $Q = \sqrt{\mathbf{q}_\perp^2} \gg \beta = 560$ MeV. *Each* of the quantities: $M_0, M'_0, \mathbf{p}_{3\perp}, \mathbf{p}_{3\perp}$ can be of order $q_\perp$, so the spin-flip term is as large as the non-spin flip term. In particular, $(s_3 = +1/2)$ may correspond to $(\lambda_3 = -1/2)$, so the spin of the struck quark $\neq$ proton spin. This means that there is no hadron helicity selection rule^{5,8}.

The effects of the lower components of Dirac spinors, which cause the spin flip term $\boldsymbol{\sigma} \times \mathbf{p}_3$, are the same as having a non-zero L_z , if the wave functions are expressed in the light-front basis. See Sect. 9.

We may now qualitatively understand the numerical results, since

$$F_1(Q^2) = \int \frac{d^2 q_\perp d\xi}{\xi(1-\xi)} \frac{d^2 K_\perp d\eta}{\eta(1-\eta)} \cdots \langle \uparrow \mathbf{p}'_3 | \uparrow(\mathbf{p}_3) \rangle \quad (17)$$

$$Q\kappa F_2(Q^2) = 2M_N \int \frac{d^2 q_\perp d\xi}{\xi(1-\xi)} \frac{d^2 K_\perp d\eta}{\eta(1-\eta)} \cdots \langle \uparrow \mathbf{p}'_3 | \downarrow(\mathbf{p}_3) \rangle, \quad (18)$$

where the $\dots$ represents common factors. The term $F_1 \sim \langle \uparrow \mathbf{p}'_3 | \uparrow \mathbf{p}_3 \rangle$ is a spin-non-flip term and $F_2 \sim \langle \uparrow \mathbf{p}'_3 | \uparrow \mathbf{p}_3 \rangle$ depends on the spin-flip term. In doing the integral each of the momenta, and M_0, M'_0 can take the large value Q for some regions of the integration. Thus in the integral

$$\langle \uparrow \mathbf{p}'_3 | \uparrow \mathbf{p}_3 \rangle \sim \frac{Q}{Q}, \quad \langle \uparrow \mathbf{p}'_3 | \downarrow \mathbf{p}_3 \rangle \sim \frac{Q}{Q}, \quad (19)$$

so that F_1 and QF_2 have the same Q^2 dependence. This is shown in Fig. 1.




Figure 1. Calculation of Refs. ^{2,5}, data are from Jones *et al.*¹⁰ and from Gayou *et al.*

9. Relation between ordinary Dirac Spinors and L_z

Our use of ordinary Dirac spinors corresponds to the use of a non-zero L_z in the light front basis. We may represent Dirac spinors as Melosh rotated Pauli spinors, and this is sufficient to show $L_z \neq 0$.

It is worthwhile to consider the pion as an explicit example. Then our version of the light-front wave function χ_π would be¹¹:

$$\chi_\pi(k^+, \mathbf{k}_\perp, \lambda, \lambda') \propto \langle \lambda | i\sigma_2 m - (k_1 - i\sigma_3 k_2) | \lambda' \rangle, \quad (20)$$

while the Gousset-Pire-Ralston⁵ pion Bethe-Salpeter amplitude Φ is

$$\Phi = P_{0\pi} \not{p}_\pi + P_{1\pi} [\not{p}_\pi, \not{k}_\perp], \quad (21)$$

where p_π is the pion total momentum, $P_{i\pi}$ are scalar functions of relative momentum, and the term with $P_{1\pi}$ is the one which carries orbital angular momentum. The relation¹² between the Bethe-Salpeter amplitude and the light-front wave function ϕ_π is

$$\phi_\pi(k^+, \mathbf{k}_\perp, \lambda, \lambda') = \bar{u}_L(k^+, \mathbf{k}_\perp, \lambda) \gamma^+ \Phi \gamma^+ v_L(P_\pi^+ - k^+, -\mathbf{k}_\perp, \lambda'). \quad (22)$$

Doing the Dirac algebra and choosing suitable functions $P_{i\pi}$, leads to $\chi_\pi = \phi_\pi$. The Melosh transformed Pauli spinors, which account for the lower components of the ordinary Dirac spinors, contain the non-zero angular momentum of the wave function Φ .

10. Neutron Charge Form Factor

The neutron has no charge, $G_{En}(Q^2 = 0) = 0$, and the square of its charge radius is determined from the low Q^2 limit as $G_{En}(Q^2) \rightarrow -Q^2 R^2/6$. The quantity R^2 is well-measured¹³ as $R^2 = -0.113 \pm 0.005 \text{ fm}^2$. The Galster parameterization¹⁴ has been used to represent the data for $Q^2 < 0.7 \text{ GeV}^2$.

Our proton respects charge symmetry, the interchange of u and d quarks, so it contains a prediction for neutron form factors. This is shown in Fig. 2. The resulting curve labeled relativistic quarks is both large and small. It is very small at low values of Q^2 . Its slope at $Q^2 = 0$ is too small by a factor of five, if one compares with the straight line. But at larger values of Q^2 the prediction is relatively large.

Our model gives $R_{\text{model}}^2 = -0.025 \text{ fm}^2$, about five times smaller than the data. The small value can be understood in terms of $F_{1,2}$. Taking the definition (2) for small values of Q^2 gives

$$-Q^2 R^2/6 = -Q^2 R_1^2/6 - \kappa_n Q^2/4M^2 = -Q^2 R_1^2/6 - Q^2 R_F^2/6, \quad (23)$$

where the Foldy contribution, $R_F^2 = 6\kappa_n/4M^2 = -0.111 \text{ fm}^2$, is in good agreement with the experimental data. That a point particle with a magnetic moment can explain the charge radius has led some to state that G_E is not a measure of the structure of the neutron. However, one must include the Q^2 dependence of F_1 which gives R_1^2 . In our model $R_1^2 = +0.086 \text{ fm}^2$ which nearly cancels the effects of R_F^2 . Isgur¹⁷ showed that this cancellation is a natural consequence of including the relativistic effects of the lower components of the Dirac spinors. Thus our relativistic effects are standard. We need another source of R^2 . This is the pion cloud.




Figure 2. Calculation of G_E^n . The data are from Ref. ¹⁵, with more expected soon¹⁶.

11. Pion Cloud and the Light Front Cloudy Bag Model

The effects of chiral symmetry require that sometimes a physical nucleon can be a bare nucleon emersed in a pion cloud. An incident photon can interact electromagnetically with a bar nucleon, a pion in flight or with a nucleon while a pion is present. These effects were included in the cloudy bag model, and are especially pronounced for the neutron. Sometimes the neutron can be a proton plus a negatively charged pion. The tail of the pion distribution extends far out into space (see Figs. 10 and 11) of Ref.¹⁸, so that the square of the charge radius is negative.

It is necessary to modernize the cloudy bag model, so as to make it relativistic. This involves using photon-nucleon form factors from our model, using a relativistic π -nucleon form factor, and treating the pionic contributions relativistically by doing a light front calculation. This has been done. The result is the light front cloudy bag model, and the preliminary results are shown in Fig. 3. We see that the pion cloud effects are important for small values of Q^2 and, when combined with those of the relativistic quarks coming from the bare nucleon, leads to a good description of the low Q^2 data. The total value of G_E is substantial for large values of Q^2 .




Figure 3. Light front cloudy bag model LFCBM Calculation of G_E^n .

12. Summary

Poincaré invariance is needed to describe the new exciting experimental results. Ordinary Dirac spinors carry light front orbital angular momentum. Including the effects of these spinors, in a way such that the proton is an eigenstate of spin leads naturally to the result that QF_2/F_1 is constant for values of Q^2 between 2 and about 20 GeV^2 .

The prediction of hadron helicity conservation is that Q^2F_2/F_1 is constant, so we see that this is not respected in present data and there is no need to expect it to hold for a variety of exclusive reactions occurring at high $Q^2 \leq 5.5 \text{ GeV}^2$. Examples include the anomalies seen in pp elastic scattering and the large spin effects seen in the reactions $\gamma d \rightarrow np$ and $\gamma p \rightarrow \pi^0 p$.

The results for the neutron G_E can be concisely stated. At small values of Q^2 the effects of a pion cloud is needed to counteract the relativistic effects which cancel the effects of the Foldy term. At large values of Q^2 relativistic effects give a “large” value of G_E ; large in the sense that this form factor is predicted to be larger than that of the Galster parameterization.

At the time of this workshop, I had not yet used the light front cloudy bag model to compute proton form factors or the neutron’s G_M . Including

the effects of the pion cloud (with a parameter to describe the pion-nucleon form factor) allows the use of different quark-model parameters. The result is an excellent description of all four nucleon electromagnetic form factors, and I plan to publish that soon.

Acknowledgments

This work is partially supported by the U.S. DOE. I thank R. Madey for encouraging me to compute the neutron form factors.

References

1. M.R. Frank, B.K. Jennings and G.A. Miller, Phys. Rev. C **54**, 920 (1996).
2. L. L. Frankfurt and M. I. Strikman, Nucl. Phys. B **250**, 143 (1985).
3. F. Schlumpf, U. Zurich Ph. D. Thesis, hep-ph/9211255.
4. G. A. Miller and M. R. Frank, nucl-th/0201021 to appear Phys. Rev. C.
5. P. Jain, B. Pire and J. P. Ralston, Phys. Rept. **271**, 67 (1996) ; T. Gousset, B. Pire and J. P. Ralston, Phys. Rev. D **53**, 1202 (1996)
6. V. B. Berestetskii and M. V. Terent'ev. Sov. J. Nucl. Phys. **25**, 347 (1977).
7. P. L. Chung and F. Coester. *Phys. Rev. D* **44**, 229, (1991).
8. V. M. Braun, A. Lenz, N. Mahnke and E. Stein, Phys. Rev. D **65**, 074011 (2002).
9. M. K. Jones *et al.*, Phys. Rev. Lett. **84**, 1398 (2000)
10. O. Gayou *et al.*, Phys. Rev. Lett. **88**, 092301 (2002).
11. P.L. Chung, F. Coester and W.N. Polyzou, Phys. Lett 205B,545 (1988).
12. H. H. Liu and D. E. Soper, Phys. Rev. D **48**, 1841 (1993).
13. S. Kopecky *et al.*, Phys. Rev. Lett. **74**, 2427 (1995)
14. S. Galster *et al.*, Nucl. Phys. B **32**, 221 (1971).
15. T. Eden *et al.*, Phys. Rev. C **50**, 1749 (1994); M. Meyerhoff *et al.*, Phys. Lett. B **327**, 201 (1994); M. Ostrick *et al.*, Phys. Rev. Lett. **83**, 276 (1999). J. Becker *et al.*, Eur. Phys. J. A **6**, 329 (1999). I. Passchier *et al.*, Phys. Rev. Lett. **82**, 4988 (1999) D. Rohe *et al.*, Phys. Rev. Lett. **83**, 4257 (1999). H. Zhu *et al.* Phys. Rev. Lett. **87**, 081801 (2001)
16. Jefferson Laboratory Experiment 93-038, R. Madey Spokesperson; R. Madey, for the Jlab E93-038 collaboration, "Neutron Electric Form Factor Via Recoil Polarimetry", contribution to Baryons 2002.
17. N. Isgur, Phys. Rev. Lett. **83**, 272 (1999)
18. S. Th  berge, A. W. Thomas and G. A. Miller, Phys. Rev. **D22** (1980) 2838; (1981) 2106, A. W. Thomas, S. Th  berge, and G. A. Miller, Phys. Rev. **D24** (1981) 216; S. Th  berge, G. A. Miller and A. W. Thomas, Can. J. Phys. **60**, 59 (1982). G. A. Miller, A. W. Thomas and S. Th  berge, Phys. Lett. B **91**, 192 (1980).